

# Tissue Stability following Soft-tissue Augmentation and Zirconia Prosthesis: A 6 Years Follow-up

Lanka Mahesh, Sangeeta Dhir, Kelvin Ian Afrashtehfar

## ABSTRACT

**Background:** 'Reconstructive esthetic implant therapy' is a term evolved in the ever-demanding age of oral implantology today. Maintaining the critical factor in this delicate balance of the gingival/pink esthetics is the periimplant papilla. Its presence being a major esthetic achievement and its loss leads to tremendous esthetic handicap known as 'black hole disease'. This case report addresses this fascinating yet challenging aspect of reconstruction of periimplant papilla through a novel technique.

**Methods:** A 38-year-old patient presented for replacement of missing tooth. Complete oral and periodontal examination, orthodontic intervention to prepare the implant site was undertaken. Preoperative soft-tissue assessments were done. A modified palatal roll technique was performed for reconstruction of the papillary architecture. Zirconia prosthesis was fabricated and the pink esthetic score (PES) around the prosthesis was calculated.

**Results:** An appreciable PES score around the prosthesis complete fill of the interproximal papilla and buccal soft tissue augmentation resulted from this technique.

**Keywords:** Papilla reconstruction, Palatal roll technique, Maxillary anterior esthetics.

**How to cite this article:** Mahesh L, Dhir S, Afrashtehfar KI. Tissue Stability following Soft-tissue Augmentation and Zirconia Prosthesis: A 6 Years Follow-up. *Int J Oral Implant Clin Res* 2012;3(2):77-82.

**Source of support:** Nil

**Conflict of interest:** None declared

## INTRODUCTION

Optimal esthetics for implant-supported restorations in the anterior maxilla may be more difficult to obtain than implant osseointegration.

The ability to predictably preserve or reproduce inter-implant papilla is extremely important in the replacement of maxillary anterior teeth. The presence of interproximal papilla around implant-supported restorations allows symmetrical soft tissues margins and a state of harmony between natural teeth and dental implant components.<sup>1</sup> This harmony and tissue symmetry leads to natural looking restoration that does not obscure vision. On the contrary, slightest change in the level of the interproximal papilla can lead to major esthetic and phonetic complications. This is what makes the periimplant supported tissues a delicate clinical issue to handle.

Ideal esthetic outcomes demands an optimal position and inclination of the implant. Periimplant soft tissue is the crucial element that decides and influences the tooth length, color, texture to impart a natural appearance to the implant-supported restoration.<sup>2</sup>

## Biologic Truths

The sequence of losing the interproximal papilla starts immediately after tooth extraction. The thin adjacent alveolar bone (interradicular bone) starts rapid resorption due to thin alveolar nature of alveolar bone, reduced blood supply to the crest of interradicular bone, possible direct contamination of the crest of interradicular bone by oral bacteria as a result of tooth extraction, absence of sharpeys fibers that stimulate continuous bone remodeling.

Engquist et al 1995<sup>3</sup> stated that consequence of tooth extraction leads to the interdental papilla remodeling in a sloping fashion from the palatal to the more apical facial osseous plate and becomes depressed in comparison with the healthy adjacent marginal tissue. Unfortunately, the lost interdental papilla cannot regenerate to regain its original dimensions.<sup>4</sup>

## Biology of the Periimplant Mucosa

There is a significant difference between the tissues surrounding the natural teeth and implants. In implants due to lack of cement-like structures, connective tissue fibers of the periimplant mucosa are stretched parallel to the implant surface rather than perpendicularly attached to the root surface as seen in natural teeth. Most groups of surracestral fibers (gingivodental and transeptal fibers) do not exist surrounding the implant abutment.

Important vital differences also is the restricted blood supply, which is due to the absence of periodontal ligament and associated blood vessel branch. The branches from the bone and oral soft tissues only provide the blood supply to the periimplant mucosa. In natural teeth, the gingival vascularization is derived from the branches originating from the interdental septa, periodontal ligament and oral mucosa. Further, the periimplant mucosa contains a high amount of collagen and low number of fibroblasts, therefore, the periimplant mucosa can also be defined as 'scar-like tissues'.

## Biologic Width around Natural Teeth and Implant

The biologic width around an implant differs from the natural tooth. Average being about 3 mm compared with that present around natural teeth about 2 mm. The junctional epithelium is also the double the size of that in healthy gingival.<sup>5</sup> Between an implant and abutment, a space or microgap always exists and biologic width forms apical to the microgap leading to crestal bone loss of about 2 mm irrespective of whether the microgap is located at or below the alveolar crest.<sup>6</sup> This indicates that crestal bone changes are not dependant on the surgical technique<sup>7</sup> (submerged/nonsubmerged) neither on the older thought of 'saucerization'<sup>8</sup> (owing to mechanical stress generated by implant body) but on the location of the interface microgap<sup>9</sup> and lack of viable biologic width around the implant.<sup>5</sup>

## FACTORS INFLUENCING THE OUTCOME OF PAPILLA RECONSTRUCTION

### Blood Supply

It's the key factor in predicting the treatment outcome, as sufficient blood supply should be maintained in any flap design.

### Implant Positioning

Well-placed implants lead to an esthetically successful implant restoration.<sup>10</sup> Three different directions govern the positioning of the implant: Apicoincisal, mesiodistal, labiopalatal.

### Periodontal Biotype

Periodontal biotype thick or thin affects the dimension of the periodontal tissue and should be carefully evaluated during presurgical planning. Thick biotype more prone to pocket formation but reconstruction procedures seem to be more predictable due to sturdy nature of the soft-tissues and osseous structure. Thin biotype more prone to gingival recession following mechanical and surgical manipulation.

### Bone Quality and Quantity

The bony support between a tooth and an implant or between two implants has been shown to be an important criterion in creating or preserving the papilla.<sup>11,12</sup> Tarnow et al reported a mean papillary height between two adjacent implants as 3.4 mm. One difficulty in maintaining or reforming a papilla between two implants is that the biologic width around an implant usually is located apically to the implant abutment junction. In the esthetic zone, the distance from the alveolar crest to the adjacent tooth CEJ should be

3 to 5 mm to achieve ideal implant localization and appropriate space for the periimplant sulcus to form.<sup>13</sup>

### Soft-tissue Quantity and Quality

The documented literature unanimously states that sufficiently broad cuff of keratinized mucosa is necessary to allow for predictable manipulation of the soft-tissue surrounding the implant gingival tissues and also leads to long-term success of oral endosseous implants and maintenance of the integrity of the interproximal papilla.<sup>14-16</sup> Of all the methods used for soft-tissue augmentation and the flap designs used, the underlying concept is to preserve the blood supply to the adjacent papilla and to minimize recession.

### Implant Size Selection

Selection of an implant for an esthetic zone depends on the dimensions of the edentulous crest and proximity of adjacent roots. Implants with larger diameter are of limited use as they compromise the interimplant distance of 3 mm leading to increased crestal bone loss.<sup>17</sup> Hence, implants 3.75 to 4 mm in diameter are preferred for the anterior restoration.<sup>18</sup> Platform switching to a smaller diameter at the interface level favors the biologic width development in the horizontal direction to compensate for vertical one hence forth minimizing the postoperative bone resorption and maintaining soft-tissue margins.<sup>19</sup>

### Emergence Profile

A proper emergence profile is important for hygiene, gingival health and appearance. Implant restorations in the esthetic zone should mimic the emergence profile (flat) of the natural tooth. The vertical length of the subgingival portion of the restoration is extremely important as the guided gingival growth is indirectly proportional to the submergence depth of the implant.<sup>20</sup> The emergence profile of the final prosthesis should be carefully created. If the profile is too narrow, no contralateral pressure or support for the gingival will exist and the interdental papilla will diminish. If the profile is too wide papilla will be vertically compressed, oral hygiene will be difficult or impossible to perform and papilla will collapse.

### Classification for Interdental Papilla

Nordland had classified according to marginal level of papilla in relation to the CEJ<sup>21</sup>:

*Class 1:* Tip of the interdental papilla lies between the interdental contact point and most coronal extent of the interproximal CEJ.

*Class 2:* Tip of the interdental papilla lies at or apical to the CEJ (interproximal CEJ visible)

*Class 3:* Tip of the interdental papilla lies level with the or apical to the facial CEJ.

Tarnow's classification had its basis on the predictability of the presence of interdental papilla:<sup>22</sup>

*Class 1:* When the distance between the contact point of the natural tooth and crest of the bone was 5 mm or less papilla is present almost 100% time

*Class 2:* When the distance was 6 mm, the papilla was present 56% times

*Class 3:* When the distance was 7 mm the papilla was present 27% times. Several other classifications have described this clinical entity.<sup>23-25</sup>

The choice of restoration for the esthetic rehabilitation in implant dentistry primarily focuses on the position, inclination, shape and color of the restoration.<sup>26</sup> The soft tissue around the implant-supported restorations play a significant role in defining its close imitation in their appearance to the natural teeth.<sup>27</sup> Restorations can cause discoloration of the mucosa.<sup>28</sup> Metal-free restorations/all ceramic restorations allow to preserve soft tissue color more similar to the natural tooth than the PFM restorations.<sup>29</sup> Zirconia with its mechanical properties similar to metal and color similar to tooth color.<sup>30</sup> Periimplant soft-tissue reaction to zirconium dioxide is minimal as compared to other restorative materials.<sup>31</sup> Biologic response of titanium vs the zirconia healing caps was studied. Results revealed increased inflammatory infiltrate, microvessel density and vascular endothelial growth factor expression around the titanium caps than around ZrO<sub>2</sub> ones.<sup>32</sup>

### The PES Scoring Criteria<sup>33</sup>

This criteria is based on the seven variables: Mesial papilla, distal papilla, soft-tissue level, soft-tissue contour, alveolar process deficiency, soft-tissue color and texture. Each variable is assessed with a score of 0-1-2, with 2 being the best and 0 being the poorest score (Table 1).



**Fig. 1:** Orthodontic treatment done to prepare the site for implant

### CASE REPORT

A 30-year-old female patient presented for routine examination with a desire of replacement of missing anterior tooth that was extracted some years ago. Patient was healthy with no significant medical history. Intraoral examination revealed congenitally missing lateral incisor in relation to left maxillary quadrant (Fig. 1). Long-standing edentulism had resulted in mesial migration of canine. Patient's oral hygiene status was adequate. Thorough treatment planning for the mouth was charted that included recording of the oral hygiene scores (plaque index, gingival bleeding index) was done. No significant periodontal findings were evident. Radiographic findings of the mouth revealed normal bone levels.

Orthodontic treatment was started with the aim of preparing the edentulous site for receiving the implant-supported prosthesis (see Fig. 1).


### PREOPERATIVE ASSESSMENTS

#### Soft-tissue Characteristics

Under anesthesia, the surgical site was sounded for the soft-tissue thickness and to record the papilla height. Mucosal thickness 4.8 mm (midbuccal), 1.75 mm (crestal) was recorded. Papilla height index 5.5 mm (Tarnow's index) was scored. Patient had a thin flat biotype in the area.

**Table 1:** PES score criteria<sup>33</sup>

Variables	Details	0	1	2
Mesial papilla	Shape vs reference tooth	Absent	Incomplete	Complete
Distal papilla	Shape vs reference tooth	Absent	Incomplete	Complete
Marginal tissue level	Level vs reference tooth	Major discrepancy >2 mm	Minor discrepancy 1 to 2 mm	No discrepancy <1 mm
Soft-tissue contour	Natural matching with reference tooth	Unnatural	Fairly natural	Natural
Alveolar process	Alveolar process deficiency	Obvious	Slight	None
Soft-tissue color	Color vs reference tooth	Obvious difference	Moderate difference	No difference
Soft-tissue texture	Texture vs reference tooth	Obvious difference	Moderate difference	No difference



**Fig. 2:** Stage 1 implant surgery force direction indicator (FDI) in position



**Fig. 4:** Flat gingival architecture at 4 months



**Fig. 3:** Implant in position



**Fig. 5:** Stage 2 implant surgery—palatal site deepithelized

**The Technique**

Stage 1 surgery was performed and 3.8/10.5 mm (tapered internal, Biohorizon, AL, USA) implant was placed following manufacturer’s protocol (Figs 2 and 3).

After 4 months Stage 2 surgery of uncovering the implant was performed along with the desired soft-tissue augmentation (Fig. 4). After securing anesthesia, esthetic flap design was planned preserving the interdental papilla along the adjacent teeth (Fig. 5). A 15-c scalpel (HU-FRIEDY, CH, USA) is used to mark the vertical incision extending from the buccal aspect toward the palatal side. The palatal extent is marked about 5 to 7 mm from the crestal tissue. Palatal site is marked and deepithelized with scalpel/diamond bur (Fig. 6). Full thickness flap is elevated from the palatal side. This elevation extends toward the buccal side as pouch dissection. Once the buccal dissection has been completed, the palatal tissue of the pedicle graft is then rolled and positioned into the buccal pouch and under the buccal flap (Fig. 7). Narrow diameter healing abutment



**Fig. 6:** Papilla preserving flap incision, flap reflected buccally

was screwed in. The pedicle graft is meticulously sutured around the emerging healing abutment using 4-0 eptfe cytoplast sutures (Osteogenics Biomedical, TX-USA). Simple interrupted sutures were placed at the mesial and



**Fig. 7:** Palatal flap rolled under the buccal flap



**Fig. 9:** Composite crown 18 months postoperative buccal view



**Fig. 8:** Temporary crown—at insertion



**Fig. 10:** Soft tissue profile after removal of composite crown

distal interproximal part of the pedicle graft that forms the future papilla and maintaining the buccal fullness simultaneously. Impression was recorded at the same visit for the fabrication of the temporary crown (Fig. 8). Two weeks postoperatively sutures were removed and temporary crown was cemented (see Fig. 8). After 8 months, with soft-tissue attaining the early stability, composite crown was placed in (Fig. 9). Two years later, impression was done to replace the composite crown with zirconia prosthesis (Figs 10 to 12).

## RESULTS

At 2 years, the composite crown was replaced with the zirconia crown. Complete fill of the interproximal papilla (mesial and distal) was observed. Gingival tissue thickness as measured on the buccal aspect was 4 mm at the crestal region and 6 mm at the mid buccal aspect. Six years follow-up has been satisfactory.



**Fig. 11:** Final abutment

## CONCLUSION

Ideal outcome of the practice of implantology depends on the functional and esthetic success. Reconstruction of the gingival esthetics is an important issue in modern esthetic implant dentistry. Ideal treatment planning and sound



**Fig. 12:** Zirconia crown in place

preoperative assessment of soft and hard tissues form the baseline for successful predictability of the reconstructed papilla. The PES scoring criteria with its advantage of reproducibility helps in assessing the success of the surgical and prosthetic protocols especially around the single tooth implant restorations.

## REFERENCES

1. Tarnow DP, Escow RN 1996. Esthetics and implant dentistry. *Periodontol* 2000;11:85-94.
2. Chang, et al. Implant supported single-tooth replacements compared to contralateral natural teeth. *Clin Oral Implan Res* 1999;10:185-94.
3. Engquist, BH Nilson, P Astrand. Single tooth replacement by osseointegrated Branemark implants. *Clin Oral Implant Res* 1995;(6):38-45.
4. Holmes CH. Morphology of the interdental papilla. *J Periodont* 1965;(36):21-26.
5. Cochran DL, Hermann JS, Schenk RK. Biologic width around titanium implants. A histometric analysis of the implantogingival junction around unloaded and loaded nonsubmerged implants in the canine mandible. *J Periodontol* 1997;68:186-98.
6. Hermann JS, et al. Crestal bone changes around titanium implants. A radiographic evaluation of unloaded nonsubmerged implants in the canine mandible. *J Periodontol* 1997;68:1117-30.
7. Weber HP, Cochran DL. Soft tissue response to osseointegrated dental implants. *J Prosthet Dent* 1998;79:79-89.
8. Mihalko WM, May TC. Finite element analysis of interface geometry effects on the crestal bone surrounding the dental implants. *Implant Dent* 1992;1(3):212-17.
9. Piaetelli A, Vrespa G. Role of the microgap between implant and abutment. *J Periodontol* 2003;(74)3:346-52.
10. Speilman HP. Influence of the implant position on the esthetics of the restoration. *Prac Periodontics Aesthet Dent* 1996;8:897-904.
11. Hartman GA, Cochran DL. Initial implant position determines the magnitude of crestal bone remodeling. *J Periodontol* 2004;75(4):572-77.
12. Choquet V, Tarnow DP. Clinical and radiographic evaluation of the papilla level adjacent to single tooth dental implants. *J Periodontol* 2001;72(10):1364-71.
13. Tarnow D, Fletcher P. Vertical distance from the crest of the bone to the height of the interproximal papilla between adjacent implants. *J Periodontol* 2003;74:1785-88.
14. Krekler G, Diemer J. Should the exit of the artificial abutment tooth be positioned in the region of the attached gingival? *Intl J Oral Surg* 1985;14:504-08.
15. Scharf DR, Tarnow DP. Modified roll technique for localized ridge augmentation. *Int J Perio Restorative Dent* 1992;7:303-10.
16. Bengazi F, Wennstorm JL. Recession of soft tissue margins in oral implants. *Clin Oral Implants Res* 1996;7:303-10.
17. Tarnow DP, Cho SC. The effect of inter-implant distance on the height of inter-implant bone crest. *J Periodontol* 2000;71:546-49.
18. Boudrias P. Evaluation of the osseous edentulous ridge- probing technique using a measuring guide. *J Dent Que* 2003;40:301-02.
19. Tarnow D. Personal Communications. NYU, USA 2004.
20. Steins JM, Nevins M. The relationship of the guided gingival frame to the provisional crown for a single implant restoration. *Compend Contin Educ Dent* 1996;17:1175-82.
21. Nordland WP, Tarnow DP. Classification system for loss of papillary height. *J Periodontol* 1998;69:1124-26.
22. Tarnow DP, Fletcher P. The effect of the distance from the contact point to the crest of the bone on the presence or absence of the interproximal papilla. *J Periodontol* 1992;63:995-96.
23. Salama H, Salama D. The interproximal height of bone—a guidepost to predictable esthetic strategies and soft tissue contours in the anterior tooth replacement. *Prac Periodontics Aesthet Dent* 1998;10:1131-41.
24. Gastaldo JF, Curyfi R. Effect of the vertical and horizontal distances between adjacent implants and between tooth and implant on the incidence of interproximal papilla. *J Periodontol* 2004;75:1242-46.
25. Lee D, Kim C. Noninvasive method to measure the length of soft tissue from the top of the papilla to the crestal bone. *J Periodontol* 2005;76:1311-14.
26. Philips K, Kois JC. Esthetic peri-implant site development. The restorative connection. *Dent Clin North Am* 1998;42:57-70.
27. Chang M, Wennstrom JL. Implant supported single tooth replacements compared to contralateral natural teeth. Crown and soft tissue dimensions. *Clin Oral Implan Res* 1999;10:185-94.
28. Takeda T, Ishigami K. A study of the discoloration of the gingival by artificial crowns. *Int J Prosthodont* 1996;9:197-202.
29. Jung RE, Sailer I, Hammerle CHF, Attin T, Schmidlin P. In vitro color changes of soft-tissues caused by restorative materials. *Int J Periodont and Restorative Dent* 2007;27:251-57.
30. Piconi C, Maccauro G. Zirconia as a ceramic biomaterial. *Biomaterials* 1999;20:1-25.
31. Warashina H, Sakano S, Kitamura S, et al. Biological reaction to alumina, zirconia, titanium and polyethylene particles implanted onto murine calvaria. *Biomaterials* 2003;24:3655-61.
32. Degidi M, et al. Inflammatory infiltrate, microvessel density, nitric oxide synthase expression, vascular endothelial growth factorexpression, and proliferative activity in peri-implant soft tissues around titanium and zirconium oxide healing caps. *J Periodontol* 2006;77:73-80.
33. Furhauser R, et al. Evaluation of soft tissue around single tooth implant crowns: The pink esthetic score. *Clin Oral Implan Res* 2005;16:639-44.

## ABOUT THE AUTHORS

### Lanka Mahesh (Corresponding Author)

Private Practice, The Dental Centre, S-382, Panshila Park New Delhi-17, India, e-mail: drlanka.mahesh@gmail.com

### Sangeeta Dhir

Professor, Department of Periodontics and Oral Implantology, Sudha Rustagi College of Dental Sciences and Research, Faridabad, Haryana, India

### Kelvin Ian Afrashtehfar

DDS, Montreal, Canada